

## Letter to the Editor

# Malignant Melanoma and Charcot-Marie-Tooth Disease: A Further Case

### To the Editor:

In a previous issue of this journal, Greene et al. [1980] described 2 patients with Charcot-Marie-Tooth (CMT) disease who later developed cutaneous malignant melanoma. Although the development of the two diseases in the same patient may have occurred by chance, the authors raised the possibility of a shared neural crest defect or a genetic linkage.

Among the patients reported by Greene et al. [1980], one had a dominant form of CMT. The patient's mother and brother were similarly affected. A paternal aunt died of melanoma. The second patient had a neuronal type of CMT. His brother showed the same disease, but the parents were not examined.

Our case involves a 29-year-old woman who came to our Clinical Genetic Unit for an assessment of her reproductive risk. Family history showed that her brother was affected by CMT [hereditary motor-sensory neuropathy (HMSN) type II] as indicated by electromyography (EMG) and sural nerve biopsy. Neurological examination of the *proposita* showed moderate reduction of the osteotendinous reflexes on the lower limbs; the EMG showed a slow nerve conduction (particularly in the right peroneal nerve). On this basis the patient was given the diagnosis of CMT (HMSN type II).

Personal history showed that she had undergone a surgical removal of a pigmented malignant melanoma from the left flank at age 26 years. Microscopically epithelioid cells were described; no invasion of the dermoepidermal junction was observed, suggesting the diagnosis of intraepidermal melanoma. Chemotherapy was not given.

The parents of our patient were normal on neurological examination and following EMG.

Charcot-Marie-Tooth disease shows clinical variability and genetic heterogeneity [Dyck and Lambert, 1968a,b]. It was proposed that the forms of CMT with very slow nerve conduction be given the gene symbols CMT1A and CMT1B CMT1A being the gene on chromosome arm 17p, and CMT1B being the gene on chromosome arm 1q. CMT2 was the proposed symbol for the autosomal locus responsible for moderately slow

nerve conduction form of the disease. The genetic types CMT1A, CMT1B, and CMT2 relate to the clinical types abbreviated HMSNIA, HMSNIB, and HMSN-II [McKusick, 1994]. The designation CMT2, or HMSN-II, should be confined to inherited axonal neuropathy (Harding, personal communication); this gene is on chromosome arm 1p.

Our case further suggests a causal association between Charcot-Marie-Tooth disease and malignant melanoma. It is interesting that, although heterogeneity is demonstrated, the most favored location of one of the CMT2 genes is on chromosome 1 within the region p35-p36 [Ben Othmane et al., 1993]; one of the cutaneous malignant melanoma genes (CMM1) also maps to chromosome band 1p36 [Dracopoli et al., 1989; Goldstein et al., 1993].

### REFERENCES

- Ben Othmane K, Middleton LT, Loprest LJ, Wilkinson KM, Lennon F, Rozear MP, Stajich JM, Gaskell PC, Roses AD, Pericak-Vance MA, Vance JM (1993): Localization of a gene (CMT2A) for autosomal dominant Charcot-Marie-Tooth disease type 2 to chromosome 1p and evidence of genetic heterogeneity. *Genomics* 17:370-375.
- Dracopoli NC, Bale SJ, Housman DE (1989): Assignment of the familial melanoma gene to chromosome 1p36: Frequent loss of heterozygosity of this region occurs late in tumor progression. *Am J Hum Genet* 45:19.
- Dyck PJ, Lambert EH (1968a): Lower motor primary sensory neuron diseases with peroneal muscular atrophy. I. Neurologic, genetic, and electrophysiologic findings in hereditary polineuropathies. *Arch Neurol* 18:603-618.
- Dyck PJ, Lambert EH (1968b): Lower motor primary sensory neuron diseases with peroneal muscular atrophy. II. Neurologic, genetic, and electrophysiologic findings in various neuronal degenerations. *Arch Neurol* 18:619-625.
- Goldstein AM, Dracopoli NC, Ho EC, Fraser MC, Kearns KS, Bale SJ, McBride OW, Clark WH, Tucker MA (1993): Further evidence for a locus for cutaneous malignant melanoma-dysplastic nevus (CMM/DN) on chromosome 1p, and evidence for genetic heterogeneity. *Am J Hum Genet* 52:537-550.
- Greene MH, Mead GD, Reimer RR, Bergfeld WF, Fraumeni JF Jr (1980): Malignant melanoma and Charcot-Marie-Tooth disease. *Am J Med Genet* 5:69-71.
- McKusick VA (1994): "Mendelian Inheritance in Man." Baltimore and London: The Johns Hopkins University Press, p 285.

**Siranoush Manoukian**  
**Vincenza Briscioli**  
**Faustina Lalatta\***  
Laboratorio di Citogenetica  
Istituti Clinici di Perfezionamento  
Milan, Italy

\*Correspondence to: Faustina Lalatta, M.D., Laboratorio di Citogenetica, Istituti Clinici di Perfezionamento, via Commenda 12, 20122 Milan, Italy.

Received 27 January 1995; Accepted 2 April 1996